



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

on the inferior thin margin of the body, and the lateral undulatory motion of the animal in the water exactly resembles that of *Cerophora*, or *Carinaria*.

The heart of *Phyllirrhoë*, in common with that of Heteropods in general, holds a dorsal position. The auricle lies posterior to the ventricle, as in *Cerophora* and *Firola*, but the reverse is the case in *Atlanta* and *Carinaria*, the difference being due to the relation which the respiratory surface bears to the heart itself, lying in every case on the auricular side. Moreover it is remarkable that the rectum is directed backwards in the former instances, but turns forwards in the latter, taking an opposite course to that of the circulation through the heart.

It may be observed in conclusion, that in Heteropoda the viscera are closely packed together so as to occupy the smallest possible space, while they are widely distributed through the abdomen in *Phyllirrhoë*; thus, again, calling to mind its relationship to the Pteropoda.

This paper is illustrated with drawings representing the animal described and some of the details of its internal structure.

IV. "Brief sketch of the Anatomy of a new genus of pelagic Gasteropoda, named *Jasonilla*." By JOHN DENIS MACDONALD, Esq., R.N. Communicated by Sir W. BURNETT, K.C.B. Received March 30, 1855.

This communication refers to a remarkable genus of pelagic Gasteropoda, characterized, like *Macgillivraya* and *Cheletropis*, by the presence of ciliated cephalic appendages, but having, as in the present instance, a beautifully transparent, cartilaginous and perfectly symmetrical shell. The author has seen but one species, which was frequently taken between Port Jackson and the Isle of Pines.

The shell resembles that of *Argonauta* in shape, is less than one-eighth of an inch in diameter, and the little animal, when fully retracted, occupies but a small portion of its cavity. The margin of the mantle is of considerable thickness, containing loosely-packed cells, similar to those of the middle or operculigerous lobe of many Pteropods. About eight ciliated arms, identical in character with those of *Macgillivraya*, &c., encircle the head, including the mouth, which

is furnished with two massive lateral jaws bearing sharp prominent dental processes on the anterior border, and with a pair of simple tentacula having a dark ocellus at the outer side of the base of each. A well-formed foot arises by a narrow pedicle from the under surface of the body, immediately behind the ciliated collar. The creeping disc is elongated in form, subquadrate in front, and tapers off gradually towards the posterior extremity. The latter part, corresponding to the operculigerous lobe of other species, is speckled with little clusters of dark pigment-cells, disposed so much after the manner of those of the ciliated arms as to lead to the impression that it is one of the same series, or whorl of organs, to use botanical phraseology. A pectinate gill extends beneath the mantle, along the anterior third of the dorsal region, lying, as in most cases, in advance of the heart. The visceral mass of the body, though elongated, is but slightly curved upon itself, not exceeding half a turn. The lobules of the liver, distended with large amber-coloured oil-globules, may be distinctly seen through the transparent outer envelope and shell. Single spherical otolithes are contained in the acoustic sacs, and the lingual ribbon is lengthy and flexuous, presenting a row of uncini on each side, with a series of minute denticulations, pointing backwards on their anterior and posterior borders. The uncini of opposite sides interlock with one another so closely as to conceal the rudimentary segments of the rachis almost completely. The shell is cartilaginous, transparent, planorbicular, and perfectly symmetrical, presenting four rows of minute conical tuberculations on its convex or dorsal surface. The gyri of the involute nucleus are so curved as to leave a central perforation; the mouth of the tube encroaches considerably on the last whorl, and the outer lip is deeply notched between the two lateral rows of tubercles. The author has named the species *Jasonilla M^cLeayiana*. The paper is accompanied with illustrative figures.

V. Note "On the position of Aluminum in the Voltaic series."

By CHARLES WHEATSTONE, Esq., F.R.S. Received April 25, 1855.

Having, through the kindness of Dr. Hofmann, been permitted to examine a specimen of aluminum prepared by M. Claire-Deville, I